

MATHEMATICS

1. If $A = \{a, b, c\}$, $B = \{b, c, d\}$ and $C = \{a, d, c\}$, then $(A - B) \times (B \cap C) =$
- 1) $\{(a, c), (a, d), (b, d)\}$ 2) $\{(c, a), (d, a)\}$
3) $\{(a, b), (c, d)\}$ 4) $\{(a, c), (a, d)\}$
2. The function $f : X \rightarrow Y$ defined by $f(x) = \sin x$ is one-one but not onto if X and Y are respectively equal to,
- 1) $\left[\frac{-\pi}{2}, \frac{\pi}{2}\right]$ and $[-1, 1]$ 2) $\left[0, \frac{\pi}{2}\right]$ and $[-1, 1]$
3) $[0, \pi]$ and $[0, 1]$ 4) \mathbb{R} and \mathbb{R}
3. If $\log_4^2 + \log_4^4 + \log_4^{16} + \log_4^x = 6$, then $x =$
- 1) 32 2) 8
3) 4 4) 64
4. If $S_n = \frac{1}{6.11} + \frac{1}{11.16} + \frac{1}{16.21} + \dots$ to n terms, then $6S_n =$
- 1) $\frac{1}{(5n+6)}$ 2) $\frac{(2n-1)}{5n+6}$
3) $\frac{n}{(5n+6)}$ 4) $\frac{5n-4}{5n+6}$
5. The remainder obtained when $(\underline{1})^2 + (\underline{2})^2 + (\underline{3})^2 + \dots + (\underline{100})^2$ is divided by 10^2 is
- 1) 14 2) 17
3) 28 4) 27

(Space for Rough Work)

6. If $(p \wedge \neg r) \rightarrow (\neg p \vee q)$ is false, then the truth values of p , q and r are respectively

- 1) T, F and T
- 2) F, T and T
- 3) F, F and T
- 4) T, F and F

7. If α , β and γ are the roots of the equation $x^3 - 8x + 8 = 0$, then $\sum \alpha^2$ and $\sum \frac{1}{\alpha \beta}$ are respectively =

- 1) 16 and 0
- 2) -16 and 0
- 3) 16 and 8
- 4) 0 and -16

8. The g.c.d. of 1080 and 675 is

- 1) 125
- 2) 225
- 3) 135
- 4) 145

9. If $a | (b+c)$ and $a | (b-c)$ where $a, b, c \in N$ then,

- 1) $c^2 \equiv a^2 \pmod{b^2}$
- 2) $a^2 \equiv b^2 \pmod{c^2}$
- 3) $a^2 + c^2 = b^2$
- 4) $b^2 \equiv c^2 \pmod{a^2}$

10. If a, b and $c \in N$ which one of the following is not true ?

- 1) $a | b$ and $a | c \Rightarrow a | b+c$
- 2) $a | b+c \Rightarrow a | b$ and $a | c$
- 3) $a | b$ and $b | c \Rightarrow a | c$
- 4) $a | b$ and $a | c \Rightarrow a | 3b+2c$

(Space for Rough Work)

11. If $2A + 3B = \begin{bmatrix} 2 & -1 & 4 \\ 3 & 2 & 5 \end{bmatrix}$ and $A + 2B = \begin{bmatrix} 5 & 0 & 3 \\ 1 & 6 & 2 \end{bmatrix}$, then $B =$

1) $\begin{bmatrix} 8 & 1 & 2 \\ 1 & 10 & 1 \end{bmatrix}$

2) $\begin{bmatrix} 8 & 1 & -2 \\ -1 & 10 & -1 \end{bmatrix}$

3) $\begin{bmatrix} 8 & 1 & 2 \\ -1 & 10 & -1 \end{bmatrix}$

4) $\begin{bmatrix} 8 & -1 & 2 \\ -1 & 10 & -1 \end{bmatrix}$

12. If $O(A) = 2 \times 3$, $O(B) = 3 \times 2$, and $O(C) = 3 \times 3$, which one of the following is not defined?

1) $C(A + B')$

2) $C(A + B')'$

3) BAC

4) $CB + A'$

13. If $A = \begin{bmatrix} 1 & -3 \\ 2 & K \end{bmatrix}$ and $A^2 - 4A + 10I = A$, then $K =$

1) 1 or 4

2) 4 and not 1

3) -4

4) 0

14. The value of $\begin{vmatrix} x+y & y+z & z+x \\ x & y & z \\ x-y & y-z & z-x \end{vmatrix} =$

1) 0

2) $(x+y+z)^3$

3) $2(x+y+z)^3$

4) $2(x+y+z)^2$

15. On the set Q of all rational numbers the operation * which is both associative and commutative is given by $a * b =$

1) $2a + 3b$

2) $ab + 1$

3) $a^2 + b^2$

4) $a + b + ab$

(Space for Rough Work)

16. In the group $G = \{1, 5, 7, 11\}$ under multiplication modulo 12, the solution of $7^{-1} \times_{12} (x \times_{12} 11) = 5$ is $x =$

 - 1) 11
 - 2) 7
 - 3) 1
 - 4) 5

17. A subset of the additive group of real numbers which is not a sub group is

 - 1) $(Q, +)$
 - 2) $(N, +)$
 - 3) $(Z, +)$
 - 4) $(\{0\}, +)$

18. If $\vec{p} = \hat{i} + \hat{j}$, $\vec{q} = 4\hat{k} - \hat{j}$ and $\vec{r} = \hat{i} + \hat{k}$, then the unit vector in the direction of $3\vec{p} + \vec{q} - 2\vec{r}$ is

 - 1) $\hat{i} + 2\hat{j} + 2\hat{k}$
 - 2) $\frac{1}{3}(\hat{i} - 2\hat{j} + 2\hat{k})$
 - 3) $\frac{1}{3}(\hat{i} - 2\hat{j} - 2\hat{k})$
 - 4) $\frac{1}{3}(\hat{i} + 2\hat{j} + 2\hat{k})$

19. If \vec{a} and \vec{b} are the two vectors such that $|\vec{a}| = 3\sqrt{3}$, $|\vec{b}| = 4$ and $|\vec{a} + \vec{b}| = \sqrt{7}$, then the angle between \vec{a} and \vec{b} is

 - 1) 150°
 - 2) 30°
 - 3) 60°
 - 4) 120°

20. If \vec{a} is vector perpendicular to both \vec{b} and \vec{c} , then

 - 1) $\vec{a} \cdot (\vec{b} \times \vec{c}) = 0$
 - 2) $\vec{a} \times (\vec{b} \times \vec{c}) = \vec{0}$
 - 3) $\vec{a} \times (\vec{b} + \vec{c}) = \vec{0}$
 - 4) $\vec{a} + (\vec{b} + \vec{c}) = \vec{0}$

(Space for Rough Work)

21. If the area of the parallelogram with \vec{a} and \vec{b} as two adjacent sides is 15 sq. units, then the area of the parallelogram having $3\vec{a} + 2\vec{b}$ and $\vec{a} + 3\vec{b}$ as two adjacent sides in sq. units is
- 1) 45 2) 75
3) 105 4) 120
22. The locus of the point which moves such that the ratio of its distances from two fixed points in the plane is always a constant $K (< 1)$ is
- 1) circle 2) straight line
3) ellipse 4) hyperbola
23. If the lines $x + 3y - 9 = 0$, $4x + by - 2 = 0$ and $2x - y - 4 = 0$ are concurrent, then $b =$
- 1) 0 2) 1
3) 5 4) -5
24. The lines represented by $ax^2 + 2hxy + by^2 = 0$ are perpendicular to each other if
- 1) $h = 0$ 2) $h^2 = ab$
3) $a + b = 0$ 4) $h^2 = a + b$
25. The equation of the circle having $x - y - 2 = 0$ and $x - y + 2 = 0$ as two tangents and $x + y = 0$ as a diameter is
- 1) $x^2 + y^2 = 1$ 2) $x^2 + y^2 = 2$
3) $x^2 + y^2 - 2x + 2y - 1 = 0$ 4) $x^2 + y^2 + 2x - 2y + 1 = 0$

(Space for Rough Work)

26. If the length of the tangent from any point on the circle $(x-3)^2 + (y+2)^2 = 5r^2$ to the circle $(x-3)^2 + (y+2)^2 = r^2$ is 16 units, then the area between the two circles in sq. units is
- 1) 16π 2) 8π
3) 4π 4) 32π
27. The circles $ax^2 + ay^2 + 2g_1x + 2f_1y + c_1 = 0$ and $bx^2 + by^2 + 2g_2x + 2f_2y + c_2 = 0$ ($a \neq 0$ and $b \neq 0$) cut orthogonally if
- 1) $g_1g_2 + f_1f_2 = c_1 + c_2$ 2) $bg_1g_2 + af_1f_2 = bc_1 + ac_2$
3) $g_1g_2 + f_1f_2 = bc_1 + ac_2$ 4) $g_1g_2 + f_1f_2 = ac_1 + bc_2$
28. The equation of the common tangent of the two touching circles, $y^2 + x^2 - 6x - 12y + 37 = 0$ and $x^2 + y^2 - 6y + 7 = 0$ is
- 1) $x + y + 5 = 0$ 2) $x + y - 5 = 0$
3) $x - y + 5 = 0$ 4) $x - y - 5 = 0$
29. The equation of the parabola with vertex at $(-1, 1)$ and focus $(2, 1)$ is
- 1) $y^2 - 2y - 12x + 13 = 0$ 2) $y^2 - 2y + 12x + 11 = 0$
3) $x^2 + 2x - 12y + 13 = 0$ 4) $y^2 - 2y - 12x - 11 = 0$
30. The equation of the line which is tangent to both the circle $x^2 + y^2 = 5$ and the parabola $y^2 = 40x$ is
- 1) $2x + y + 5 = 0$ 2) $2x - y - 5 = 0$
3) $2x - y + 5 = 0$ 4) $2x - y \pm 5 = 0$

(Space for Rough Work)

31. $x = 4(1 + \cos \theta)$ and $y = 3(1 + \sin \theta)$ are the parametric equations of

1) $\frac{(x-4)^2}{16} + \frac{(y-3)^2}{9} = 1$

2) $\frac{(x-4)^2}{16} - \frac{(y-3)^2}{9} = 1$

3) $\frac{(x+4)^2}{16} + \frac{(y+3)^2}{9} = 1$

4) $\frac{(x-3)^2}{9} + \frac{(y-4)^2}{16} = 1$

32. If the distance between the foci and the distance between the directrices of the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$
 are in the ratio $3 : 2$, then $a : b$ is =

1) $2 : 1$

2) $1 : 2$

3) $\sqrt{3} : \sqrt{2}$

4) $\sqrt{2} : 1$

33. The ellipse $\frac{x^2}{25} + \frac{y^2}{16} = 1$ and the hyperbola $\frac{x^2}{25} - \frac{y^2}{16} = 1$ have in common

1) centre and vertices only

2) centre, foci and vertices

3) centre, foci and directrices

4) centre only

34. If $\sec \theta = m$ and $\tan \theta = n$, then $\frac{1}{m} \left[(m+n) + \frac{1}{(m+n)} \right] =$

1) mn

2) $2n$

3) $2m$

4) 2

35. The value of $\frac{\sin 85^\circ - \sin 15^\circ}{\cos 65^\circ} =$

1) 0

2) 1

3) -1

4) 2

(Space for Rough Work)

36. From an aeroplane flying vertically above a horizontal road, the angles of depression of two consecutive stones on the same side of the aeroplane are observed to be 30° and 60° respectively. The height at which the aeroplane is flying in km is

1) 2

$$2) \quad \frac{2}{\sqrt{3}}$$

$$3) \frac{\sqrt{3}}{2}$$

$$4) \quad \frac{4}{\sqrt{3}}$$

37. If the angles of a triangle are in the ratio $3 : 4 : 5$, then the sides are in the ratio

1) 3 : 4 : 5

$$2) \quad 2 : \sqrt{3} : \sqrt{3} + 1$$

$$3) \quad \sqrt{2} : \sqrt{6} : \sqrt{3} + 1$$

$$4) \quad 2 : \sqrt{6} : \sqrt{3} + 1$$

38. If $\cos^{-1}x = \alpha$, $(0 < x < 1)$ and $\sin^{-1}\left(2x\sqrt{1-x^2}\right) + \sec^{-1}\left(\frac{1}{2x^2-1}\right) = \frac{2\pi}{3}$,

then $\tan^{-1}(2x) =$

$$1) \quad \frac{\pi}{2}$$

$$2) \quad \frac{\pi}{3}$$

$$3) \quad \frac{\pi}{4}$$

$$4) \frac{\pi}{6}$$

- 39.** If $a > b > 0$, then the value of $\tan^{-1}\left(\frac{a}{b}\right) + \tan^{-1}\left(\frac{a+b}{a-b}\right)$ depends on

1) neither a nor b

2) a and not b

3) b and not a

4) both a and b

- 40.** Which one of the following equations has no solution?

$$1) \quad \sqrt{3} \sin \theta - \cos \theta = 2$$

$$2) \quad \cos \theta + \sin \theta = \sqrt{2}$$

$$3) \quad \operatorname{Cosec} \theta \cdot \operatorname{Sec} \theta = 1$$

$$4) \quad \operatorname{Cosec} \theta = \operatorname{Sec} \theta \equiv \operatorname{Cosec} \theta : \operatorname{Sec} \theta$$

(Space for Rough Work)

41. The complex number $\frac{(-\sqrt{3} + 3i)(1-i)}{(3+\sqrt{3}i)(i)(\sqrt{3} + \sqrt{3}i)}$ when represented in the Argand diagram lies
- 1) on the X-axis (Real axis)
 - 2) on the Y-axis (Imaginary axis)
 - 3) in the first quadrant
 - 4) in the second quadrant
42. If $2x = -1 + \sqrt{3}i$, then the value of $(1-x^2+x)^6 - (1-x+x^2)^6 =$
- 1) 0
 - 2) 64
 - 3) -64
 - 4) 32
43. The modulus and amplitude of $(1+i\sqrt{3})^8$ are respectively
- 1) 256 and $\frac{8\pi}{3}$
 - 2) 2 and $\frac{2\pi}{3}$
 - 3) 256 and $\frac{2\pi}{3}$
 - 4) 256 and $\frac{\pi}{3}$
44. The value of $\lim_{x \rightarrow 0} \frac{5^x - 5^{-x}}{2x} =$
- 1) $2 \log 5$
 - 2) 1
 - 3) 0
 - 4) $\log 5$
45. Which one of the following is not true always?
- 1) If a function $f(x)$ is continuous at $x=a$, then $\lim_{x \rightarrow a} f(x)$ exists.
 - 2) If $f(x)$ and $g(x)$ are differentiable at $x=a$, then $f(x)+g(x)$ is also differentiable at $x=a$.
 - 3) If $f(x)$ is continuous at $x=a$, then it is differentiable at $x=a$.
 - 4) If $f(x)$ is not continuous at $x=a$, then it is not differentiable at $x=a$.

(Space for Rough Work)

46. If $y = 1 + \frac{1}{x} + \frac{1}{x^2} + \frac{1}{x^3} + \dots \text{ to } \infty$ with $|x| > 1$, then $\frac{dy}{dx} =$

1) $\frac{-y^2}{x^2}$

2) $\frac{y^2}{x^2}$

3) $x^2 y^2$

4) $\frac{x^2}{y^2}$

47. If $f(x)$ and $g(x)$ are two functions with $g(x) = x - \frac{1}{x}$ and $fog(x) = x^3 - \frac{1}{x^3}$, then $f'(x) =$

1) $3x^2 + \frac{3}{x^4}$

2) $1 + \frac{1}{x^2}$

3) $x^2 - \frac{1}{x^2}$

4) $3x^2 - 3$

48. The derivative of $a^{\sec x}$ w.r.t. $a^{\tan x}$ ($a > 0$) is

1) $a^{\sec x - \tan x}$

2) $\sin x a^{\sec x - \tan x}$

3) $\sin x a^{\tan x - \sec x}$

4) $\sec x a^{\sec x - \tan x}$

49. If $\sin(x+y) + \cos(x+y) = \log(x+y)$, then $\frac{d^2y}{dx^2} =$

1) 1

2) -1

3) 0

4) $\frac{-y}{x}$

50. If $f(x)$ is a function such that $f''(x) + f(x) = 0$ and $g(x) = [f(x)]^2 + [f'(x)]^2$ and $g(3) = 8$, then $g(8) =$

1) 8

2) 3

3) 0

4) 5

(Space for Rough Work)

51. If the curve $y = 2x^3 + ax^2 + bx + c$ passes through the origin and the tangents drawn to it at $x = -1$ and $x = 2$ are parallel to the X-axis, then the values of a , b and c are respectively.

- 1) 3, -12 and 0 2) -3, 12 and 0
3) -3, -12 and 0 4) 12, -3 and 0

52. A circular sector of perimeter 60 metre with maximum area is to be constructed. The radius of the circular arc in metre must be

- 1) 10 2) 15
3) 5 4) 20

53. The tangent and the normal drawn to the curve $y = x^2 - x + 4$ at $P(1, 4)$ cut the X-axis at A and B respectively. If the length of the subtangent drawn to the curve at P is equal to the length of the subnormal, then the area of the triangle PAB in sq. units is

- 1) 16 2) 8
3) 32 4) 4

54. $\int \frac{(x^3 + 3x^2 + 3x + 1)}{(x+1)^5} dx =$

- 1) $\tan^{-1} x + c$ 2) $\log(x+1) + c$
3) $\frac{1}{5} \log(x+1) + c$ 4) $-\frac{1}{(x+1)} + c$

55. $\int \frac{\text{Cosec } x}{\cos^2 \left(1 + \log \tan \frac{x}{2}\right)} dx =$

- 1) $-\tan \left[1 + \log \tan \frac{x}{2}\right] + c$ 2) $\sec^2 \left[1 + \log \tan \frac{x}{2}\right] + c$
3) $\tan \left[1 + \log \tan \frac{x}{2}\right] + c$ 4) $\sin^2 \left[1 + \log \tan \frac{x}{2}\right] + c$

(Space for Rough Work)

56. $\int \frac{dx}{x\sqrt{x^6 - 16}} =$

1) $\text{Sec}^{-1}\left(\frac{x^3}{4}\right) + c$

2) $\frac{1}{12} \text{Sec}^{-1}\left(\frac{x^3}{4}\right) + c$

3) $\text{Cosh}^{-1}\left(\frac{x^3}{4}\right) + c$

4) $\frac{1}{3} \text{Sec}^{-1}\left(\frac{x^3}{4}\right) + c$

57. If $I_1 = \int_0^{\pi/2} x \sin x \, dx$ and $I_2 = \int_0^{\pi/2} x \cos x \, dx$, then which one of the following is true ?

1) $I_1 = I_2$

2) $I_1 + I_2 = 0$

3) $I_1 = \frac{\pi}{2} I_2$

4) $I_1 + I_2 = \frac{\pi}{2}$

58. If $f(x)$ is defined in $[-2, 2]$ by $f(x) = 4x^2 - 3x + 1$ and $g(x) = \frac{f(-x) - f(x)}{(x^2 + 3)}$, then

$\int_{-2}^2 g(x) \, dx =$

1) 24

2) 0

3) -48

4) 64

59. The area enclosed between the parabola $y = x^2 - x + 2$ and the line $y = x + 2$ in sq. units =

1) $\frac{4}{3}$

2) $\frac{2}{3}$

3) $\frac{1}{3}$

4) $\frac{8}{3}$

60. The solution of the differential equation $e^{-x}(y+1) \, dy + (\cos^2 x - \sin 2x)y \, (dx) = 0$ subjected to the condition that $y = 1$ when $x = 0$ is

1) $(y+1) + e^x \cos^2 x = 2$

2) $y + \log y = e^x \cos^2 x$

3) $\log(y+1) + e^x \cos^2 x = 1$

4) $y + \log y + e^x \cos^2 x = 2$

(Space for Rough Work)